



## 1 Euler's Method Basics

The main idea of Euler's method is that we are given a first-order ordinary differential equation of the form:

$$\frac{dy}{dt} = f(t, y), \quad y(t_0) = y_0$$

and we want to calculate an *approximate* solution of this equation.

With Euler's method, we start at the point  $(t_0, y_0)$  and want to approximate the value of  $y(t)$  at some time  $t_1$ . The approach of Euler's method is to approximate the solution as a straight line with slope given by  $f(t_0, y_0)$ . Then we figure out our approximate value  $y_1$  by calculating the value of this line at  $t_1$ . Algebraically, this process would be:

$$y_1 = y_0 + (t_1 - t_0)f(t_0, y_0)$$

We can then repeat this process for more steps, using  $(y_1, t_1)$  to calculate  $(y_2, t_2)$ , then using that to calculate  $(y_3, t_3)$  and so on. So more generally we have:

$$y_{n+1} = y_n + (t_{n+1} - t_n)f(t_n, y_n)$$

Often the timestep  $t_{n+1} - t_n$  is uniform, and so we call it  $h$ , in which case we have:

$$y_{n+1} = y_n + hf(t_n, y_n)$$

## 2 Euler's Method Example

Today I want to go over a basic example with Euler's method that I think is particularly instructive.

Let's consider the example of:

$$\frac{dy}{dt} = -1000y, \quad y(0) = 1$$

By now, you'll probably recognize that the exact solution is  $y(t) = e^{-1000t}$ . This function starts out at 1, and very quickly approaches 0 as  $t$  increases. We might expect that Euler's method will then give us a sequence of straight lines that resemble this curve, but let's see what actually happens.

Let's use a "small" stepsize of  $h = 0.01$ , then our Euler's method formula will become:

$$y_{n+1} = y_n + 0.01f(t_n, y_n) = y_n + 0.01(-1000y_n) = y_n - 10y_n = -9y_n$$

We start with  $y_0 = y(0) = 1$ . We calculate the first few steps:

$$\begin{aligned} y(0.01) &\approx y_1 = -9y_0 = (-9)(1) = -9 \\ y(0.02) &\approx y_2 = -9y_1 = (-9)(-9) = 81 \\ y(0.03) &\approx y_3 = -9y_2 = (-9)(81) = -729 \end{aligned}$$

And so on. The exact solution is supposed to go to 0 very quickly, but this approximation keeps switching sign and is *increasing* exponentially instead of decreasing exponentially. Why is this going on? Why is Euler's method not "working" here?

The answer is that we are not using a small enough timestep  $h$ . I picked this equation out because it is what is known as a *stiff* equation, basically meaning that it requires a very small timestep  $h$  to be computed accurately. It turns out that if we use an even smaller value of  $h$  then the approximate solution will behave correctly.

The moral of the story here is that Euler's method "works" in the sense that it converges to the correct solution as  $h \rightarrow 0$ . But that doesn't mean it is necessarily accurate for a given  $h$ . The ideal way to use numerical methods such as Euler's method is using what's often called "grid refinement": using a series of decreasing values of  $h$  and checking that the solution is converging as  $h \rightarrow 0$ .